

## Diversity of Tree Vegetation in Mount Baung Natural Tourism Park Utilization Block, Pasuruan Regency

Muhammad Hasan Ilyasa<sup>1</sup>, Dwi Suheriyanto<sup>1\*</sup>

<sup>1</sup> Biology Department, Faculty of Science and Technology, Maulana Malik Ibrahim State Islamic University of Malang, Jl. Gajayana No. 50, Malang City, East Java, Indonesia 65144.

\*Email: [dsuheriyanto@bio.uin-malang.ac.id](mailto:dsuheriyanto@bio.uin-malang.ac.id)

**ABSTRACT.** Natural forests have a wide variety of tree species. The high diversity of tree species helps the forest maintain ecological balance. Mount Baung Natural Tourism Park is the conservation area in Pasuruan Regency, East Java that functions to protect biodiversity. The research aims to identify tree species, analyze tree diversity, and determine tree dominance with the Importance Value Index (IVI). The research was conducted from February to March 2023 in Mount Baung Natural Tourism Park Utilization Block (MBNTP UB), Pasuruan Regency, East Java, Indonesia. The observation location was determined to be 37 points and plots were made in the point with a size of 20 m × 20 m for the tree phase, 10 m × 10 m for pole phase, 5 m × 5 m for the sapling phase, and 2 m × 2 m for the seedling phase. Data collected from field observations were analyzed to determine the Shannon diversity index using PAST 4.01. Tree dominance was analyzed using IVI. The research showed that 14 species from 8 tree families are found in MBNTP UB. The Fabaceae family is found to be abundant compared to other families. The highest number of individuals found in the seedling phase is *Cassia glauca*. *Swietenia macrophylla* has the highest individuals number in the sapling, pole, and tree phases. The highest tree diversity is found in the tree phase. Diversity at the tree phase is classified as moderate, while at the seedling, pole, and sapling phase the diversity is low.

**Keywords:** conservation; Fabaceae; IVI; Mount Baung; Shannon index

### INTRODUCTION

One of Indonesia's biological wealth is tree diversity. Trees are plants with woody stems and can live for years (Dulamsuren *et al.*, 2014). Trees are biotic components that can form vegetation and are dominant in a forest ecosystem, which have their respective roles (Carteron *et al.*, 2020). Forests are an essential habitat for biodiversity and provide various ecosystem services for human welfare (Brockerhoff *et al.*, 2017). Trees in forests have an essential role in avoiding erosion, regulating the hydrological cycle, maintaining global climate stability, and storing carbon. They can absorb carbon dioxide produced from various activities carried out by humans and other living things (Aba *et al.*, 2017). Trees can also eliminate air pollution through the interception of particulates on plant surfaces and the absorption of pollutant gases through leaf stomata (Nowak *et al.*, 2014).

Natural forests have a wide variety of tree species. The high diversity of tree species helps the forest maintain ecological balance (Safe'i *et al.*, 2018). Forest health status can be determined based on indicators of tree species diversity. The level of tree species diversity is closely related to an ecosystem's ecological stability level. The higher the number of tree species and the value of tree species diversity in a forest, the greater the complexity of the forest ecosystem (Sanjaya *et al.*, 2021). Tree diversity assessment is vital because it is sensitive to changes, indicators of ecological systems, spatial-temporal heterogeneity, and parts of the food chain (Safe'i *et al.*, 2021). Tree vegetation analysis activities help to understand the type and arrangement of trees in forest areas, support reforestation initiatives, and evaluate and develop forest resources (Cahyanto *et al.*, 2014).

Mount Baung Natural Tourism Park (MBNTP) is the conservation area in Pasuruan Regency, East Java, that functions to protect biodiversity. Biodiversity research in MBNTP that has ever been carried out shows that various types of plant that grow in MBNTP, including bamboo, cemberit, saga, Javanese sengon, and taro (Furqoni *et al.*, 2023). There are six bamboo species that grow with a clumped distribution pattern. This area's bamboo population is generally young phase (Sofiah *et al.*, 2013). *Syzygium* species were found in MBNTP, six species were found growing in this place. *Syzygium pycnanthum* is the most commonly found in MBNTP (Mudiana, 2016). In the MBNTP


there is a native Indonesian plant, namely *Lepisanthes rubiginosa*. The presence of *L. rubiginosa* in MBNTP is quite good and has the potential to be utilized (Mudiana & Ariyanti, 2021).

Managers and local communities have done reforestation to support MBNTP sustainability. Forest loss is the main cause of species extinction, and reforestation is one way to restore forests (Kemppinen *et al.*, 2020). Large-scale reforestation has an essential impact on plant species diversity, which in turn affects the stability and resilience of vegetation (Wang *et al.*, 2019). Reforestation can significantly improve species composition, density, and canopy cover (Osuri *et al.*, 2019). Reforestation is the primary strategy for mitigating climate change due to its potential carbon storage impacts (Locatelli *et al.*, 2015).

Information about the diversity of tree vegetation in the Mount Baung Natural Tourism Park, especially in the Utilization Block (MBNTP UB), still needs to be improved, so research needs to be done. The research aims to identify tree species, analyze tree diversity, and determine tree dominance with the Importance Value Index (IVI). The research results can provide an overview of the structure and composition of tree species in MBNTP UB Pasuruan Regency, making it helpful in determining management strategies for conservation or restoration.

## MATERIALS AND METHODS

**Study area.** The research was conducted from February to March 2023 in MBNTP UB, Pasuruan Regency, East Java, Indonesia is located at  $112^{\circ}16'23''$  -  $112^{\circ}17'17''$  E and  $07^{\circ}46'09''$  -  $07^{\circ}47'23''$  S, at altitude 282 m asl. Environmental conditions varied widely, with temperatures around 28 - 32 °C, humidity around 51 - 60%, wind speeds around 1.5 - 9 m/s, and light intensity around 1766 - 6572 lux.



**Fig. 1.** Study area in the Mount Baung Natural Tourism Park Utilization Block.

**Procedures.** The vegetation analysis sampling begins with determining the research location point in the MBNTP UB area. Based on the survey results, the area that can be accessed is 15 ha. Sampling was carried out on 10% of the area using a quadratic plot. Plots were made in the point with a size of  $20 \times 20$  m for the tree phase,  $10 \times 10$  m for pole phase observations,  $5 \times 5$  m for the sapling phase, and  $2 \times 2$  m for the seedling phase. The number of sample points in the MBNTP UB is 37 (Fig. 1). Observations were made and recorded regarding the type and number of trees in each plot and

morphological observations of each fruit, seed, flower, and leaf. Then, the morphology of each fruit, seed, flower, and leaf sample was identified using Steenis (2005) and Sukarya (2013).

**Data analysis.** Data collected from field observations were analyzed to determine the Shannon diversity index using PAST 4.01. Shannon diversity index criteria: <1.5: low, 1.5-3.5: moderate, >3.5: high (Magurran, 1988). Tree dominance was analyzed using the Important Value Index (IVI) (Fachrul, 2007). The IVI formula is as follows:

Density (D):

$$D = \frac{\text{Number of Individuals}}{\text{total area sampled}}$$

Relative Density (RD):

$$RD = \frac{\text{Density of a species}}{\text{Total density for all species}} \times 100\%$$

Frequency (F):

$$F = \frac{\text{Number of plots in which species recorded}}{\text{Total number of plots sampled}}$$

Relative Frequency (RF):

$$RF = \frac{\text{Frequency of a species}}{\text{Total frequency for all species}} \times 100\%$$

Dominance (Dom):

$$\text{Dom} = \frac{\text{Total basal area of a species}}{\text{Total area sampled}}$$

Relative Dominance (RDom):

$$\text{RDom} = \frac{\text{Dominance of a species}}{\text{Total dominance for all species}} \times 100\%$$

IVI formula for seedling and sapling:

$$\text{IVI} = \text{RD} + \text{RF}$$

IVI formula for pole and tree:

$$\text{IVI} = \text{RD} + \text{RF} + \text{RDom}$$

## RESULTS AND DISCUSSION

### The Tree Identification

Based on research results, it is known that there are 14 species from 8 tree families found in MBNTP UB. The families found are Anacardiaceae, Euphorbiaceae, Fabaceae, Malvaceae, Meliaceae, Moraceae, Sapindaceae, and Verbenaceae (Table 1).

**Table 1.** The tree identification results in Mount Baung Natural Tourism Park Utilization Block.

Family	Species	Local Name
Anacardiaceae	<i>Lannea coromandelica</i> (Houtt.) Merr.	Mentaos
Euphorbiaceae	<i>Mallotus molissimus</i> (Geiseler) Airy Shaw	Tutup
Fabaceae	<i>Samanea saman</i> (Jacq.) Merr.	Trembesi
Fabaceae	<i>Delonix regia</i> (Bojer ex Hook.) Raf.	Flamboyan
Fabaceae	<i>Cassia glauca</i> Lam.	Joar
Fabaceae	<i>Parkia timoriana</i> (DC.) Merr.	Kedawung
Fabaceae	<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	Sengon buto
Fabaceae	<i>Butea monosperma</i> (Lam.) Taubert	Ploso
Fabaceae	<i>Vachellia nilotica</i> (L.) P.J.H.Hurter & Mabb.	Klampis
Malvaceae	<i>Ceiba pentandra</i> (L.) Gaertn.	Randu
Meliaceae	<i>Swietenia macrophylla</i> King.	Mahoni
Moraceae	<i>Ficus racemosa</i> L.	Elo
Sapindaceae	<i>Sapindus rarak</i> Dc.	Klerek
Verbenaceae	<i>Tectona grandis</i> L.f.	Jati

The Fabaceae family is found to be abundant compared to other families. There are 7 species of the Fabaceae family found in MBNTP UB. This family has much potential to be used. The Fabaceae family is an essential group of plants because humans use it as legumes, green manure, and forage. This plant also has many benefits in pharmacology (Ahmad *et al.*, 2016). There are 23 genera from the Fabaceae family used in Bangladesh to treat various skin, respiratory, circulatory, digestive, and various diseases (Rahman & Parvin, 2014). In Zimbabwe, 101 species of the Fabaceae family are used for medicine. These plants are used as traditional medicine to treat 134 medical conditions related to respiratory, digestive, female reproductive, and sexually transmitted infections (Maroyi, 2023).

### The Tree Diversity

Based on Table 2, The three highest numbers in the seedling phase are *C. glauca* (146 individuals), *S. macrophylla* (130 individuals), and *S. rarak* (66 individuals). In the sapling phase are *S. macrophylla* (121 individuals), *C. glauca* (106 individuals), and *S. rarak* (34 individuals). In the pole phase are *S. macrophylla* (71 individuals), *C. glauca* (55 individuals), and *S. rarak* (17 individuals). In the tree phase are *S. macrophylla* (112 individuals), *C. glauca* (44 individuals), and *T. grandis* (16 individuals). The seedling phase (385 individuals) is the highest individual number, followed by the sapling, tree, and pole with individual numbers 278, 252, and 157, respectively. The highest number of species is found in the tree phase, with 14 species, followed by the seedling phase, with 6 species, while the pole and sapling phase has the same number of species, namely 5. The highest Shannon diversity index value is in the tree phase (1.902), followed by the seedling phase (1.402), pole phase (1.219), and sapling phase (1.192). Diversity at the tree phase is classified as moderate, while at the seedling, pole, and sapling phase the diversity is low.

**Table 2.** The tree diversity in the Mount Baung Natural Tourism Park Utilization Block.

Species	Seedling	Sapling	Pole	Tree
<i>Lannea coromandelica</i> (Houtt.) Merr.				3
<i>Mallotus molissimus</i> (Geiseler) Airy Shaw		4	2	5
<i>Samanea saman</i> (Jacq.) Merr.				2
<i>Delonix regia</i> (Bojer ex Hook.) Raf.				8
<i>Cassia glauca</i> Lam.	146*	106	55	44
<i>Parkia timoriana</i> (DC.) Merr.	11	13	12	14
<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	15			6
<i>Butea monosperma</i> (Lam.) Taubert				5
<i>Vachellia nilotica</i> (L.) P.J.H.Hurter & Mabb.				2
<i>Ceiba pentandra</i> (L.) Gaertn.				6
<i>Swietenia macrophylla</i> King.	130	121*	71*	112*
<i>Ficus racemosa</i> L.				14
<i>Sapindus rarak</i> Dc.	66	34	17	15
<i>Tectona grandis</i> L.f.	17			16
Individual number	385*	278	157	252
Species number	6	5	5	14*
Shannon index	1.402	1.192	1.219	1.902*

Note: \* : the highest value

The research results show that *S. macrophylla* has the highest numbers in the pole, sapling, and tree phases. *S. macrophylla* is widespread in various forest types, from savanna to tropical rainforest, and is often found as a border tree in mixed hardwood forests and along river banks (Telrandhe *et al.*, 2022). Adaptation capacity influences the species survival. *S. macrophylla* can live well in a variety of environmental conditions. They can tolerate various soils and is resistant to strong winds because it has strong roots (Krisnawati *et al.*, 2011). The ability to survive *S. macrophylla* in unsuitable conditions is good. This species is relatively tolerant to low rainfall and can survive in areas with



sufficient soil moisture (Brown *et al.*, 2003). The annual growth of *S. macrophylla* is susceptible to climate and is positively correlated with rainfall in the first six years (Susatya & Yansen, 2016).

Seedling dynamics can influence forest diversity. The high seed mortality rate and slow seedling growth make it difficult for trees to survive. The high mortality rate in the early seedling phase significantly limits seedling growth to the sapling phase. Seedlings with a fast and consistent growth rate can develop into trees (Chang-Yang *et al.*, 2021). The occurrence of dynamic changes in the species number, density, and dominance of plants from the seedling phase to the tree phase indicates a reciprocal relationship in vegetation that is beneficial or otherwise. One of the reciprocal relationships in plant communities can be shown by the competition of individual plants between species and within the species itself. Therefore, vegetation is defined as a whole consisting of a combination of various plant species that depend on their environment and influence each other (Mueller & Ellenberg, 2016).

The high diversity of tree phases compared to other phases shows that MBNTP UB as a conservation area has been established for a long time. The tree diversity level is directly proportional to the resilience level. The higher tree diversity level in a forest, the greater the resilience level of the forest (Safe'i *et al.*, 2018). Tree diversity is the primary driver of biodiversity in forests, the functional composition of trees, forest structure, climate, and soil. Mixed forests have many taxa, resulting in high tree diversity (Ampoorter *et al.*, 2020).

### **The Tree Dominance**

Based on Table 3, it is known that the seedling phase is dominated by *C. glauca* (IVI = 69.27%). The sapling, pole, and tree phase is dominated by *S. macrophylla* with IVI = 81.53%, 134.51%, and 112.13%, respectively. One of the factors that caused the large number of *C. glauca* species found in the seedling phase was the characteristics of *C. glauca*, a plant with a good level of adaptation to the environmental conditions of MBNTP UB. The presence of seeds different types in the same individual of *C. glauca* has the potential to grow in colonies, thus requiring different germination requirements in various environmental conditions (Khurana & Singh, 2001). As an adaptive species, *C. glauca* can survive in relatively dry conditions. This species can overcome drought by reducing water loss through stomata on its leaves or by developing strong roots to absorb water more efficiently (Putri & Dharmono, 2018).

The tree widely planted by managers and local communities for the conservation in MBNTP UB is *S. macrophylla*, so the plant is dominant in the pole, sapling, and tree phases. The presence of *S. macrophylla* can improve soil conditions, and has great potential for reforestation and greening. This plant is also used in agroforestry systems with corn, upland rice, and cassava (Krisnawati *et al.*, 2011). Research shows that *S. macrophylla* was found to be the most dominant in the Dhaka South City Corporation Urban Area, Bangladesh, with an IVI of 193.22% (Jaman *et al.*, 2017). Other researchers found *S. macrophylla* was also dominant in the seedling, sapling, pole, and tree phases in Nglanggeran Village of Batur Agung zone, Gunungkidul District, Yogyakarta (Tohirin *et al.*, 2021).

Forest managers have the most significant influence on the growth rate of *S. macrophylla* after harvest by implementing reforestation (Free *et al.*, 2014). The main goal of reforestation is to restore ecosystem functions and increase species diversity (Derhé *et al.*, 2016). Selecting tree species for reforestation programs in species-rich forest ecosystems requires much consideration to achieve landscape reforestation goals (Chechina & Hamann, 2015). The reforestation process needs to consider the types of trees planted. Choosing various tree types is more important than simply increasing the number of certain tree types (Jactel *et al.*, 2017).

The existence of trees in MBNTP UB has excellent potential to be utilized. The high potential of the Natural Tourism Park area requires protection, preservation, and sustainable use to maintain its diversity (Kurey *et al.*, 2019). Sustainable forest management has implemented various frameworks to maintain biodiversity (Mori *et al.*, 2016). Trees have implications for planning structures,

management institutions, and governance so that they can be used to improve sustainability, adaptation, and mitigation efforts (Ellison *et al.*, 2017).

**Table 3.** Important Value Index of phases in the tree.

Phase	Species	RD (%)	RF (%)	RDom (%)	IVI (%)
Seedling	<i>Cassia glauca</i> Lam.	37.92	31.34		69.27*
	<i>Parkia timoriana</i> (DC.) Merr.	2.86	5.97		8.83
	<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	3.90	5.97		9.87
	<i>Swietenia macrophylla</i> King.	33.77	28.36		62.12
	<i>Sapindus rarak</i> Dc.	17.14	20.90		38.04
	<i>Tectona grandis</i> L.f.	4.42	7.46		11.88
Sapling	<i>Mallotus molissimus</i> (Geiseler) Airy Shaw	1.44	2.00		3.44
	<i>Cassia glauca</i> Lam.	38.13	42.00		80.13
	<i>Parkia timoriana</i> (DC.) Merr.	4.68	6.00		10.68
	<i>Swietenia macrophylla</i> King.	43.53	38.00		81.53*
	<i>Sapindus rarak</i> Dc.	12.23	12.00		24.23
Pole	<i>Mallotus molissimus</i> (Geiseler) Airy Shaw	1.27	1.47	1.24	3.98
	<i>Cassia glauca</i> Lam.	35.03	33.82	33.30	102.15
	<i>Parkia timoriana</i> (DC.) Merr.	7.64	13.24	5.56	26.44
	<i>Swietenia macrophylla</i> King.	45.22	41.18	48.11	134.51*
	<i>Sapindus rarak</i> Dc.	10.83	10.29	11.79	32.91
Tree	<i>Lannea coromandelica</i> (Houtt.) Merr.	1.20	2.63	0.46	4.29
	<i>Mallotus molissimus</i> (Geiseler) Airy Shaw	2.01	3.51	0.80	6.32
	<i>Samanea saman</i> (Jacq.) Merr.	0.80	0.88	3.94	5.62
	<i>Delonix regia</i> (Bojer ex Hook.) Raf.	3.21	5.26	3.99	12.46
	<i>Cassia glauca</i> Lam.	17.67	20.18	7.35	45.19
	<i>Parkia timoriana</i> (DC.) Merr.	5.62	8.77	7.23	21.62
	<i>Enterolobium cyclocarpum</i> (Jacq.) Griseb.	2.41	4.39	6.71	13.50
	<i>Butea monosperma</i> (Lam.) Taubert	2.01	1.75	5.63	9.39
	<i>Vachellia nilotica</i> (L.) P.J.H.Hurter & Mabb.	0.80	0.88	1.24	2.92
	<i>Ceiba pentandra</i> (L.) Gaertn.	2.41	3.51	8.64	14.55
	<i>Swietenia macrophylla</i> King.	44.98	25.44	41.71	112.13*
	<i>Ficus racemosa</i> L.	5.62	10.53	2.05	18.20
	<i>Sapindus rarak</i> Dc.	6.02	9.65	4.28	19.95
<i>Tectona grandis</i> L.f.	6.43	5.26	6.44	18.13	

Note: \* : the highest value

## CONCLUSION

There are 14 species from 8 tree families found in MBNTP UB. The Fabaceae family is found to be abundant compared to other families. The highest number of individuals found in the seedling phase is *C. glauca*. Species has the highest individual numbers in the sapling, pole, and tree phases is *S. macrophylla*. The highest tree diversity is found in the tree phase (1.902), followed by the seedling phase (1.402), pole phase (1.219), and sapling phase (1.192). Diversity at the tree phase is classified as moderate, while at the seedling, pole, and sapling phase the diversity is low.

## ACKNOWLEDGEMENTS

We thank to Mount Baung Natural Tourism Park and the Natural Resources Conservation Center of East Java Province for providing any information and references. We grateful to persons who are willing to help in the field work: Luthfinia Farah Dina, Farhani Nurshafa Rahmania, Anisa Kurnia, Mochammad Faa'is Murtadho, Achmad Ady Pramuja, Lilis Nurhalimah, Annisa Safari Putri, Wildan Zainuri.

## REFERENCES

- Aba SC, Ndukwe OO, Amu CJ, Baiyeri KP. 2017. The role of trees and plantation agriculture in mitigating global climate change. *African Journal of Food, Agriculture, Nutrition and Development*. vol 17(4): 12691–12707. doi: <https://doi.org/10.18697/ajfand.80.15500>.
- Ahmad F, Anwar F, Hira S. 2016. Review on medicinal importance of fabaceae family. *PharmacologyOnLine*. vol 3: 151-156. [https://pharmacologyonline.silae.it/files/archives/2016/vol3/PhOL\\_2016\\_3\\_A023\\_47\\_Ahmad.pdf](https://pharmacologyonline.silae.it/files/archives/2016/vol3/PhOL_2016_3_A023_47_Ahmad.pdf).
- Ampoorter E, Barbaro L, Jactel H, Baeten L, Boberg J, Carnol M, Castagneyrol B, Charbonnier Y, Dawud SM, Deconchat M, Smedt PD, Wandeler HD, Guyot V, Hättenschwiler S, Joly F-X, Koricheva J, Milligan H, Muys B, Nguyen D, Ratcliffe S, Raulund-Rasmussen K, Scherer-Lorenzen M, van der Plas F, Keer JV, Verheyen K, Vesterdal L, Allan E. 2020. Tree diversity is key for promoting the diversity and abundance of forest-associated taxa in Europe. *Oikos*. vol 129: 133-146. doi: <https://doi.org/10.1111/oik.06290>.
- Brockerhoff EG, Barbaro L, Castagneyrol B, Forrester DI, Gardiner B, J González-Olabarria JR, Lyver POB, Meurisse N, Oxbrough A, Taki H, Thompson ID, van der Plas F, Jactel H. 2017. Forest biodiversity, ecosystem functioning and the provision of ecosystem services. *Biodivers Conserv*. vol 26: 3005-3035. doi: <https://doi.org/10.1007/s10531-017-1453-2>.
- Brown N, Jennings S, Clements T. 2003. The ecology, silviculture and biogeography of mahogany (*Swietenia macrophylla*): a critical review of the evidence. *Perspectives in Plant Ecology, Evolution and Systematics*. vol 6(1-2): 37-49. doi: <https://doi.org/10.1078/1433-8319-00041>.
- Cahyanto T, Chairunnisa D, Sudjarwo T. 2014. Analisis vegetasi pohon Hutan Alam Gunung Manglayang Kabupaten Bandung. *Jurnal Istek*. vol 8(2): 145 – 161. <https://journal.uinsgd.ac.id/index.php/istek/article/view/225/240>.
- Carteron A, Parasquive V, Blanchard F, Guilbeault-Mayers X, Turner BL, Vellend M, Laliberté E. 2020. Soil abiotic and biotic properties constrain the establishment of a dominant temperate tree into boreal forests. *Journal of Ecology*. vol 108(3): 931–944. doi: <https://doi.org/10.1111/1365-2745.13326>.
- Chang-Yang C-H, Needham J, Lu C-L, Hsieh C-F, Sun I-F, McMahon SM. 2021. Closing the life cycle of forest trees: The difficult dynamics of seedling-to-sapling transitions in a subtropical rainforest. *J Ecol*. vol 109: 2705-2716. doi: <https://doi.org/10.1111/1365-2745.13677>.
- Chechina M, Hamann A. 2015. Choosing species for reforestation in diverse forest communities: social preference versus ecological suitability. *Ecosphere*. vol 6(11): 1-13. doi: <http://dx.doi.org/10.1890/ES15-00131.1>.
- Derhé MA, Murphy H, Monteith G, Menéndez R. 2016. Measuring the success of reforestation for restoring biodiversity and ecosystem functioning. *J Appl Ecol*. vol 53: 1714-1724. doi: <https://doi.org/10.1111/1365-2664.12728>.
- Dulamsuren C, Khishigiargal M, Leuschner C, Hauck M. 2014. Response of tree-ring width to climate warming and selective logging in Larch Forests of the Mongolian Altai. *Journal of Plant Ecology*. vol 7(1): 24-38. doi: <https://doi.org/10.1093/jpe/rtt019>.
- Ellison D, Morris CE, Locatelli B, Sheil D, Cohen J, Murdiyarso D, Gutierrez V, van Noordwijk M, Creed IF, Pokorny J, Gaveau D, Spracklen DV, Tobella AB, Ilstedt U, Teuling AJ, Gebrehiwot SG, Sands DC, Muys B, Verbist B, Springgay E, Sugandi Y, Ullivan CA. 2017. Trees, forests and water: cool insights for a hot world. *Global Environmental Change*. vol 43: 51-61. doi: <https://doi.org/10.1016/j.gloenvcha.2017.01.002>.
- Fachrul MF. 2007. Metode sampling bioekologi. Jakarta: Bumi Aksara.
- Fatima O, Seher N, Mushtaq Z. 2022. Partial purification of bioactive fractions from cassia glauca lam leaves. *Journal of Animal & Plant Sciences*. vol 32(2): 589-595. doi: <https://doi.org/10.36899/JAPS.2022.2.0458>.
- Free CM, Matthew LR, Grogan J, Schulze MD, Lentini M, Dünisch O. 2014. Management implications of long-term tree growth and mortality rates: A modeling study of big-leaf mahogany (*Swietenia macrophylla*) in the Brazilian Amazon. *Forest Ecology and Management*, vol 33: 46-54. doi: <https://doi.org/10.1016/j.foreco.2014.05.057>.
- Furqoni MI, Astina IK, Insani N. 2023. Analisis kelayakan potensi objek daya tarik wisata alam di Gunung Baung Desa Cowek Kecamatan Purwodadi Kabupaten Pasuruan. *Media Komunikasi Geografi*. vol 24(1): 45-54. doi: <https://doi.org/10.23887/mkg.v24i1.58611>.
- Jactel H, Bauhus J, Boberg J, Bonal D, Castagneyrol B, Gardiner B, Gonzalez-Olabarria JR, Koricheva J, Meurisse N, Brockerhoff EG. 2017. Tree diversity drives forest stand resistance to natural disturbances. *Curr Forestry Rep*. vol 3: 223-243. doi: <https://doi.org/10.1007/s40725-017-0064-1>.
- Jaman MS, Jahan I, Jamil M, Helal MGJ, Islam MS, Azad MJ, Hossain MF. 2017. Structure and composition of plant vegetation in urban area of Dhaka South City Corporation, Bangladesh. *Preprints*. 1-21. doi: <https://doi.org/10.20944/preprints201711.0136.v1>.
- Kemppinen KMS, Collins PM, Hole DG, Wolf C, Ripple WJ, Gerber LR. 2020. Global reforestation and biodiversity conservation. *Conservation Biology*. vol 34: 1221-1228. doi: <https://doi.org/10.1111/cobi.13478>.
- Khurana E, Singh JS. 2001. Ecology of seed and seedling growth for conservation and restoration of tropical dry forest : a review. *Environmental Conservation*. vol 28(1): 39-52. doi: <https://doi.org/10.1017/S0376892901000042>.
- Krisnawati H, Kallio MH, Kanninen M. 2011. *Swietenia macrophylla: ecology, silviculture and productivity*. Bogor: Center for International Forestry Research. p 1-24.
- Kurey AJ, Yohanes Y, Rahawarin YY, Renwarin DMH. 2019. Potensi jasa lingkungan pada kawasan Taman Wisata Alam Klamono, kabupaten Sorong. *Jurnal Kehutanan Pappusia*. vol 5(2): 124-133. doi: <https://doi.org/10.46703/jurnalpappusia.Vol5.Iss2.143>.

- Locatelli B, Catterall CP, Imbach P, Kumar C, Lasco R, Marín-Spiotta E, Mercer B, Powers JS, Schwartz N, Uriarte M. 2015. Tropical reforestation and climate change: beyond carbon. *Restor Ecol.* vol 23: 337-343. doi: <https://doi.org/10.1111/rec.12209>.
- Magurran AE. 1988. Ecological diversity and its measurement. Princeton: Princeton University Press.
- Maroyi A. 2023. Medicinal uses of the fabaceae family in Zimbabwe: a review. *Plants.* vol 12(1255): 1-26. doi: <https://doi.org/10.3390/plants12061255>.
- Mori AS, Kenneth P, Lertzman KP, Gustafsson L. 2016. Biodiversity and ecosystem services in forest ecosystems: a research agenda for applied forest ecology. *Journal of Applied Ecology.* vol 54: 12-27. doi: <https://doi.org/10.1111/1365-2664.12669>.
- Mudiana D. 2016. *Syzygium* diversity in Gunung Baug, East Java, Indonesia. *Diversity.* vol 17(2): 733-740. doi: <https://doi.org/10.13057/biodiv/d170248>.
- Mudiana D, Ariyanti EE. 2021. Katilayu (*Lepisanthes rubiginosa* (Roxb.) Leenh.) population in Mt. Baug Nature Tourism Park. *IOP Conference Series: Earth and Environmental Science.* vol 743(1): 1-14. doi: <https://doi.org/10.1088/1755-1315/743/1/012023>.
- Mueller-Dombois D, Ellenberg H. 2016. Ekologi vegetasi: tujuan dan metode. Penerjemah: Kuswata Kartawinata dan Rochadi Abdulhadi. Jakarta: LIPI Press.
- Nowak DJ, Hirabayashi S, Bodine A, Greenfield E. 2014. Tree and forest effects on air quality and human health in the United States. 2014. *Environmental Pollution.* vol 193: 119-129. doi: <https://doi.org/10.1016/j.envpol.2014.05.028>.
- Osuri AM, Kasinathan S, Siddhartha MK, Mudappa D, Raman TRS. 2019. Effects of restoration on tree communities and carbon storage in rainforest fragments of the Western Ghats, India. *Ecosphere.* vol 10(9):1-13. <https://doi.org/10.1002/ecs2.2860>.
- Perea R, Schroeder JW, Dirzo R. 2022. The herbaceous understory plant community in the context of the overstory: an overlooked component of tropical diversity. *Diversity.* vol 14(10): 1-10. doi: <https://doi.org/10.3390/d14100800>.
- Putri AI, Dharmono. 2018. Keanekaragaman genus tumbuhan dari famili fabaceae di kawasan hutan Pantai Tabanio, Kabupaten Tanah Laut, Kalimantan Selatan. *Prosiding Seminar Nasional Lingkungan Lahan Basah.* vol 3(1): 209-213. <https://snllb.ulm.ac.id/prosiding/index.php/snllb-lit/article/view/45/45>.
- Rahman AHMM, Parvin MIA. 2014. Study of medicinal uses on fabaceae family at rajshahi, Bangladesh. *Research in Plant Sciences.* vol 2(1): 6-8. <http://pubs.sciepub.com/plant/2/1/2/>.
- Safe'i R, Erly H, Wulandari C, Kaskoyo H. 2018. Analisis keanekaragaman jenis pohon sebagai salah satu indikator kesehatan hutan konservasi. *Jurnal Perennial.* vol 14(2): 32-36. doi: <https://doi.org/10.24259/v14i2.5195>.
- Safe'i R, Kaskoyo H, Darmawan A, Haikal FF. 2021. Keanekaragaman jenis pohon sebagai salah satu indikator kesehatan hutan lindung (studi kasus di kawasan hutan lindung yang dikelola oleh HKm Beringin Jaya). *Jurnal Belantara.* vol 4(1): 89-97. doi: <https://doi.org/10.29303/jbl.v4i1.601>.
- Sanjaya FA, Safe'i R, Winarno GD. 2021. Keanekaragaman jenis pohon sebagai salah satu indikator kesehatan hutan konservasi. *Indonesian Journal of Conservation.* vol 10(2): 53-57. doi: <https://doi.org/10.15294/ijc.v10i2.28895>.
- Sofiah S, Setiadi D, Widyatmoko D. 2013. Pola penyebaran, kelimpahan dan asosiasi bambu pada komunitas tumbuhan di Taman Wisata Alam Gunung Baug Jawa Timur. *Berita Biologi.* vol 12(2): 239-247. [https://e-journal.biologi.lipi.go.id/index.php/berita\\_biologi/article/view/538](https://e-journal.biologi.lipi.go.id/index.php/berita_biologi/article/view/538).
- Steenis Van CGGJ. 2005. Flora. Bandung: PT. Pradnya Paramita.
- Sukarya DG. 2013. *3500 Plant Species of The Botanic Gardens of Indonesia.* Jakarta: PT Sukarya & Sukarya Pandetama.
- Susatya A, Yansen. 2016. Dendrochronology of young *Swietenia macrophylla* and the variation of its growth response to the past wet climate in Bengkulu, Indonesia. *Biodiversitas.* vol 17: 466-472. doi: <https://doi.org/10.13057/biodiv/d170210>.
- Telrandhe UB, Kosalge SB, Parihar S, Sharma D, Hemalatha S. 2022. Collection and cultivation of *Swietenia macrophylla* King. *Sch Acad J Pharm.* vol 11(1): 13-19. doi: <https://doi.org/10.36347/sajp.2022.v11i01.003>.
- Tohirin, Suryanto P, Sadono R. 2021. Vegetation structure, aboveground biomass, and carbon storage of *wono*, local forest management in Gunungkidul, Yogyakarta, Indonesia, across three geomorphological zones. *Biodiversitas.* vol 22: 3207-3218. doi: <https://doi.org/10.13057/biodiv/d220814>.
- Wang J, Zhao W, Zhang X, Liu Y, Wang S, Liu Y, 2019. Effects of reforestation on plant species diversity on the Loess Plateau of China: A case study in Danangou catchment. *Science of The Total Environment.* vol 651(1): 979-989. doi: <https://doi.org/10.1016/j.scitotenv.2018.09.266>.