

PAPER • OPEN ACCESS

## 2018 WMA: Applied Mathematics, Computer Science, Information Systems, and Information Technology

To cite this article: 2018 *J. Phys.: Conf. Ser.* **1114** 011001

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

### Forewords from the Head of Committee

Assalamu Alaikum Warahmatullahi Wabarakatuh.

The honourable, President of the Indonesian Publication Collaboration Community (KO2PI) and all Advisors, Vice Presidents, Secretaries and Treasurers and all invited Speakers and Participants.

First of all, let us praise and thank Allah Subhana Wata'ala, for His blessings and help, we can gather here to attend this Workshop.

Secondly, I would like to welcome all delegates, speakers and participants who attended today at the Workshop on Multidisciplinary and Its Applications “Article Writing Strategy for Internationally Indexed Multidisciplinary Publication” Part Applied Mathematics, Computer Science, Information Systems and Information Technology of course with its Education, this activity will be held in Banda Aceh on June 28, 2018, organized by the Indonesian Publication Collaboration Community (KO2PI).

Thank you very much for attention.  
Assalamu Alaikum Warahmatullahi Wabarakatuh.

Regard,

Dr. Dahlan Abdullah, M.Kom.  
Chairman



### **Forewords from President of KO2PI**

Honorable Rector of UNIMAL Aceh  
Honorable Keynote Speakers  
Honorable Head of Committee  
Distinguished all speakers and guests  
All participants,  
Ladies & Gentlement,

Assalamu'alaikum Warahmatullahi Wabarakatuh

My greetings for all of you. May peace and God's blessing be upon us all.  
*Alhamdulillah*, all praises be to the Almighty God, Allah *subhanahu wata'ala*.

It is my pleasure to welcome you all to the opening of Workshop on Multidisciplinary and Its Applications "Article Writing Strategy for Internationally Indexed Multidisciplinary Publication" Part Applied Mathematics, Computer Science, Information Systems and Information Technology.

I do hope that this workshop would bring a great opportunity for all of us to strengthen our contribution to the advancement of our nation.

I would like to take this opportunity to thank the organizing committee for their diligent work. I would also like to thank participants, especially those of you coming from others province, for joining us and sharing your valuable experiences. Should you find any inconveniences and shortcomings, please accept our sincere apologies.

Finally, let me wish you fruitful discussion and a very pleasant stay in Aceh.

Thank you,  
Wassalamu'alaykum Warahmatullahi Wabarakatuh

President of KO2PI

Ansari Saleh Ahmar

## WORKSHOP SCHEDULE

**2018 WMA Mathcomtech**  
**Indonesian Publication Collaboration Community (KO2PI)**  
 Banda Aceh, Indonesia, 28 June 2018

<b>Time</b>	<b>Activities</b>	<b>Place</b>
07.00 – 08.00	Registration	Grand Nanggroe Hall
08.00 – 09.00	Opening Ceremony Traditional Dance Do'a National Anthem: Indonesia Raya Welcome speech: a. President KO2PI Mr. Ansari Saleh Ahmar b. Rector UNIMAL Prof. Dr. Apridar, SE, M.Si	Grand Nanggroe Hall
09.00 – 09.45	Keynote Speaker 1 Robbi Rahim (Similarity and Paraphrase)	Grand Nanggroe Hall
09.45 – 10.00	Coffee Break	Grand Nanggroe Hall
10.00 – 12.00	Keynote Speaker 2 Ansari Saleh Ahmar (Strategy of Publication)	Grand Nanggroe Hall
12.00 – 13.00	Lunch & Pray	
13.00 – 16.00	Keynote Speaker 3 Darmawan Napitupulu (Mendeley)	Grand Nanggroe Hall
16.00 – 16.30	Pray	
16.30 – 17.00	Closing Ceremony	Grand Nanggroe Hall

### Workshop Session



PAPER • OPEN ACCESS

## On the distance spectrum and distance energy of complement of subgroup graphs of dihedral group

To cite this article: Abdussakir *et al* 2018 *J. Phys.: Conf. Ser.* **1114** 012109

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection—download the first chapter of every title for free.

# On the distance spectrum and distance energy of complement of subgroup graphs of dihedral group

Abdussakir<sup>1</sup>, E Susanti<sup>1</sup>, Turmudi<sup>1</sup>, M N Jauhari<sup>2</sup> and N M Ulya<sup>2</sup>

<sup>1</sup> Department of Mathematics Education, Graduate Program, Universitas Islam Negeri Maulana Malik Ibrahim Malang, Jl. Gajayana 50, Malang 65144, Indonesia

<sup>2</sup> Department of Mathematics, Faculty of Science and Technology, Universitas Islam Negeri Maulana Malik Ibrahim Malang, Jl. Gajayana 50, Malang 65144, Indonesia

sakir@mat.uin-malang.ac.id

**Abstract.** Let  $G$  is a connected simple graph and  $V(G) = \{v_1, v_2, \dots, v_p\}$  is vertex set of  $G$ . The distance matrix of  $G$  is a matrix  $D(G) = [d_{ij}]$  of order  $p$  where  $[d_{ij}] = d(v_i, v_j)$  is distance between  $v_i$  and  $v_j$  in  $G$ . The set of all eigenvalues of matrix  $D(G)$  together with their corresponding multiplicities is named the distance spectrum of  $G$  and denoted by  $spec_D(G)$ . The distance energy of  $G$  is  $E_D(G) = \sum_{i=1}^p |\lambda_i|$ , where  $\lambda_i$  are eigenvalues of  $D(G)$ . In the recent paper, the distance spectrum and distance energy of complement of subgroup graphs of dihedral group are determined.

## 1. Introduction

The study on spectrum of a graph begins with the work of Bigg [1] and defined on the adjacency matrix of graph. Next developed concept of spectrum based on various matrices obtained from a graph, such as detour spectrum [2], Laplacian spectrum [3,4], signless Laplacian spectrum [5,6], distance spectrum [7], detour distance Laplacian spectrum [8] and color signless Laplacian spectrum [9]. Until now the topic of distance spectrum is still extensively studied by researchers, see [10-21].

The concept that is very close to the concept of spectrum of a graph is energy. The concept of energy of graph was first proposed by Gutman [22]. Adjacency energy is the first concept of energy that is discussed and has many benefits in the field of chemistry [12]. Further developed other concepts of energy that widely used in graphs, such as incidence energy [23], Laplacian incidence energy [24], maximum degree energy [25], matching energy [26], Harary energy [27], color energy [28] and distance energy [29]. Distance energy was first proposed by Indulal and Gutman [7,29]. Research on the distance energy of a connected simple graph also got a great attention from the researchers, see [13,30-38].

The most recent development in graph theory is the topic of graph associated with a group. Several concepts of graph obtained from a group have been introduced, for example Cayley graph [39], identity graph [40], commuting graph [41], conjugate graph [42], non commuting graph [43], subgroup graph [44] and inverse graph [45] of a group. For given normal subgroup  $H$  of a finite group  $G$ , the subgroup graph  $\Gamma_H(G)$  of  $G$  is defined as simple graph with  $V(\Gamma_H(G)) = G$  and  $xy \in E(\Gamma_H(G))$  if and only if  $xy \in H$  [43,45].



From its introduction in 2012 by Anderson et al [44], research on subgroup graph is still rare and some of them are [45,46]. So, research on the distance spectrum and energy of subgroup graph is an interesting topic to be conducted.

## 2. Literature Review

Graph in the present paper is simple and finite. Graph  $G$  is connected if there exist  $u$ - $v$  path in  $G$ , for any two vertices  $u$  and  $v$  in  $G$  [48]. Graph  $\bar{G}$  with  $V(\bar{G}) = V(G)$  and  $xy \in E(\bar{G})$  if and only if  $xy \notin E(G)$  is named as complement of  $G$  [49]. Graph  $G$  is complete if for any two distinct vertices  $x, y \in V(G)$  then  $xy \in E(G)$ . If  $V(G)$  can be partitioned into  $k > 1$  partition sets such that an edge of  $G$  joining two vertices in the different partition then  $G$  is said to be  $k$ -partite graph. A  $k$ -partite graph  $G$  is called to be complete if for every vertices in different partition set are adjacent [50]. If  $V_1, V_2, V_3, \dots, V_k$  are partition set of a complete  $k$ -partite graph and  $|V_i| = n_i$  then this graph is denoted by  $K_{n_1, n_2, n_3, \dots, n_k}$ .

Let graph  $G$  is connected with  $V(G) = \{v_1, v_2, \dots, v_p\}$  as its vertex set. The distance matrix of  $G$  is a  $p \times p$  matrix  $D(G) = [d_{ij}]$  such that  $[d_{ij}] = d(v_i, v_j)$ , where  $d(v_i, v_j)$  is the length of the shortest  $v_i$ - $v_j$  path in graph  $G$ . The characteristics polynomial of  $D(G)$  is  $p_D(\lambda) = \det(D(G) - \lambda I)$  and the roots of the characteristics equation are named eigenvalues of  $D(G)$  or simply  $D$ -eigenvalues of graph  $G$ . The set of all  $D$ -eigenvalues of  $G$  together with their multiplicities is named the distance spectrum or simply  $D$ -spectrum of  $G$  and notated by  $spec_D(G)$  [11]. Since  $D$ -matrix is real and symmetric then all  $D$ -eigenvalues are also real [34]. Let  $\lambda_1 > \lambda_2 > \dots > \lambda_k$  are distinct  $D$ -eigenvalues of  $G$  with multiplicities  $m_1, m_2, \dots, m_k$ . According to the ordinary spectrum notation [51], then  $D$ -spectrum of  $G$  is notated by

$$spec_D(G) = \begin{bmatrix} \lambda_1 & \lambda_2 & \dots & \lambda_k \\ m_1 & m_2 & \dots & m_k \end{bmatrix}.$$

The distance energy or simply  $D$ -energy  $E_D(G)$  of connected graph  $G$  is defined as

$$E_D(G) = \sum_{i=1}^p |\lambda_i| = \sum_{j=1}^k m_j |\lambda_j|.$$

Some previous results on  $D$ -spectrum and  $D$ -energy of a connected graphs that will useful for further discussion as the followings.

**Theorem 1.1.** ([51], Theorem 4.1.)

Let  $K_{n_1, n_2, n_3, \dots, n_k}$  is complete  $k$ -partite graph of order  $n = \sum_{i=1}^k n_i$ . Then, characteristic polynomial of  $D(K_{n_1, n_2, n_3, \dots, n_k})$  is

$$p_D(\lambda) = (\lambda + 2)^{n-k} \left[ \prod_{i=1}^k (\lambda - n_i + 2) - \sum_{i=1}^k n_i \prod_{j=1, j \neq i}^k (\lambda - n_j + 2) \right].$$

The next result was first conjectured by [52] and then proved by [53].

**Theorem 1.2.**

If  $n_1, n_2, \dots, n_k > 2$  then  $E_D(K_{n_1, n_2, n_3, \dots, n_k}) = 4(n_1 + n_2 + \dots + n_k - k)$ .

Based on Theorem 1.1. the following corollaries are obvious.

**Corollary 1.3.** ([12], Corollary 3.)

Distance spectrum of complete bipartite graph  $K_{n_1, n_2}$  is

$$spec_D(K_{n_1, n_2}) = \begin{bmatrix} n_1 + n_2 - 2 + \sqrt{n_1^2 - n_1 n_2 - n_2^2} & n_1 + n_2 - 2 + \sqrt{n_1^2 - n_1 n_2 - n_2^2} & -2 \\ 1 & 1 & n_1 + n_2 - 2 \end{bmatrix}$$

**Corollary 1.4.** ([12], Corollary 4.)



For  $n_1, n_2 > 2$ ,  $E_D(K_{n_1, n_2}) = 4(n_1 + n_2 - 2)$ .

### 3. Main Results

Here, we focused on the study of subgroup graphs of dihedral group  $D_{2m}$  of order  $2m$ . It was well known that for odd  $n$ , the normal subgroup of  $D_{2m}$  are  $\langle 1 \rangle$ ,  $\langle r^d \rangle$  where  $d|m$  and  $D_{2n}$  itself and for even  $m$ , the normal subgroups of  $D_{2m}$  are  $\langle 1 \rangle$ ,  $\langle r^d \rangle$  where  $d|m$ ,  $\langle r^2, s \rangle$ ,  $\langle r^2, rs \rangle$  and  $D_{2m}$  itself. Because the concept of  $D$ -spectrum and  $D$ -energy of graph are only for connected graphs, we focused on the complement of these subgroup graphs when these subgroup graphs are not connected. The results of this study as follows.

#### Theorem 3.1.

If  $m > 3$  then

$$(a) \text{spec}_D(\overline{\Gamma_{\langle r \rangle}(D_{2m})}) = \begin{bmatrix} 3m-2 & m-2 & -2 \\ 1 & 1 & 2(m-1) \end{bmatrix}.$$

$$(b) E_D(\overline{\Gamma_{\langle r \rangle}(D_{2m})}) = 8(m-1).$$

Proof.

(a) Since  $\Gamma_{\langle r \rangle}(D_{2m}) = 2K_m$ , we have  $\overline{\Gamma_{\langle r \rangle}(D_{2m})} = K_{m,m}$ . By Corollary 1.3. it is obvious that

$$\text{spec}_D(\overline{\Gamma_{\langle r \rangle}(D_{2m})}) = \begin{bmatrix} 3m-2 & m-2 & -2 \\ 1 & 1 & 2(m-1) \end{bmatrix}.$$

(b) Based on the proof of (a) and because  $m > 3$ , by Corollary 1.4. it is complete the proof.

#### Theorem 3.2.

If  $m > 3$  and  $m$  is even then

$$(a) \text{spec}_D(\overline{\Gamma_{\langle r^2 \rangle}(D_{2m})}) = \begin{bmatrix} \frac{5m-4}{2} & \frac{m-4}{2} & -2 \\ 1 & 3 & 2(m-2) \end{bmatrix}.$$

$$(b) E_D(\overline{\Gamma_{\langle r^2 \rangle}(D_{2m})}) = 8(m-2).$$

Proof.

(a) For even  $m > 3$ ,  $\Gamma_{\langle r^2 \rangle}(D_{2m}) = 4K_{m/2}$ . So, we obtain its complement is  $\overline{\Gamma_{\langle r^2 \rangle}(D_{2m})} = K_{m/2, m/2, m/2, m/2}$ . Applying Theorem 1.1. and performing some computation we have

$$p_D(\lambda) = (\lambda + 2)^{2(m-2)} \left( \lambda - \frac{5m-4}{2} \right) \left( \lambda - \frac{m-4}{2} \right)^3.$$

The  $D$ -eigenvalues of  $\overline{\Gamma_{\langle r^2 \rangle}(D_{2m})}$  are  $(5m-4)/2$ ,  $(m-4)/2$  and  $-2$  and their multiplicities are 1, 3 and  $2(m-2)$ . Then,

$$\text{spec}_D(\overline{\Gamma_{\langle r^2 \rangle}(D_{2m})}) = \begin{bmatrix} \frac{5m-4}{2} & \frac{m-4}{2} & -2 \\ 1 & 3 & 2m-4 \end{bmatrix}.$$

(b) From (a), then  $E_D(\overline{\Gamma_{\langle r^2 \rangle}(D_{2m})}) = \left| \frac{5m-4}{2} \right| + 3 \left| \frac{m-4}{2} \right| + 2(m-2)|-2| = 8(m-2)$ . We can also apply Theorem 1.2. when  $m > 4$  to get  $E_D(\overline{\Gamma_{\langle r^2 \rangle}(D_{2m})}) = 4 \left( 4 \frac{m}{2} - 4 \right) = 8(m-2)$ .

#### Theorem 3.3.

If  $m > 3$  and  $m$  is even then

$$(a) \text{spec}_D(\overline{\Gamma_{\langle r^2, s \rangle}(D_{2m})}) = \begin{bmatrix} 3m-2 & m-2 & -2 \\ 1 & 1 & 2(m-1) \end{bmatrix}.$$

$$(b) E_D(\overline{\Gamma_{\langle r^2, s \rangle}(D_{2m})}) = 8(m-1).$$

Proof.

(a) Since  $\Gamma_{\langle r^2, s \rangle}(D_{2m}) = K_m \cup K_m$ , we have  $\overline{\Gamma_{\langle r^2, s \rangle}(D_{2m})} = K_{m,m}$ . By Corollary 1.3. the proof is clear.

(b) Based on the proof of (a) and because  $m > 3$ , by Corollary 1.4. the proof is obvious.

#### Theorem 3.4.

If  $m > 3$  and  $m$  is even then

$$(a) \operatorname{spec}_D(\overline{\Gamma_{\langle r^2, rs \rangle}(D_{2m})}) = \begin{bmatrix} 3m-2 & m-2 & -2 \\ 1 & 1 & 2(m-1) \end{bmatrix}.$$

$$(b) E_D(\overline{\Gamma_{\langle r^2, rs \rangle}(D_{2m})}) = 8(m-1).$$

Proof.

(a) Since  $\Gamma_{\langle r^2, rs \rangle}(D_{2m}) = K_m \cup K_m$ , we have  $\overline{\Gamma_{\langle r^2, rs \rangle}(D_{2m})} = K_{m,m}$ . By Corollary 1.3. we prove the part (a).

(b) Based on the proof of (a) and because  $m > 3$ , by Corollary 1.4. the proof is clear.

#### 4. Conclusions

We have computed distance spectrum and energy of complement of several subgroup graphs of dihedral group  $D_{2m}$ . The remaining subgroup graphs of dihedral group  $D_{2m}$  are given to the reader for further investigation.

#### References

- [1] Biggs N. 1974. *Algebraic graph theory*. Cambridge: Cambridge Univeristy Press.
- [2] Ayyaswamy SK and Balachandran S. 2010. On detour spectra of some graphs. *Int. J. Math. Comput. Phys. Electr. Comput. Eng.* **4** 1038–40.
- [3] Newman MW. 2000. *The Laplacian Spectrum of Graphs*. Winnipeg: The University of Manitoba.
- [4] Abdussakir, Elvierayani RR and Nafisah M. 2017. On the spectra of commuting and non commuting graph on dihedral group. *Cauchy-Jurnal Mat. Murni dan Apl.* **4** 176–82.
- [5] Yu G, Wu Y and Shu J. 2011. Signless Laplacian spectral radii of graphs with given chromatic number. *Linear Algebra Appl.* **435** 1813–22.
- [6] Abdussakir and Khasanah R. 2018. Spektrum signless-Laplace dan spektrum detour graf konjugasi dari grup dihedral. *J. Kubik.* **3** 45–51.
- [7] Indulal G and Gutman I. 2008. On the distance spectra of some graphs. *Math. Commun.* **13** 123–31.
- [8] Kaladevi V and Abinayaa A. 2017. On detour distance Laplacian energy. *J. Informatics Math. Sci.* **9** 721–32.
- [9] Bhat PG and D'Souza S. 2017. Color signless Laplacian energy of graphs. *AKCE Int. J. Graphs Comb.* **14** 142–8.
- [10] Gopalapillai I. 2009. Sharp bounds on distance spectral radius and distance energy of graphs. *Linear Algebr. Its Appl. Multilinear Algebr.* **430** 106–13.
- [11] Indulal G. 2009. Distance spectrum of graph compositions. *Ars Math. Contemp.* **2**:93–100.
- [12] Stevanović D and Indulal G. 2009. The distance spectrum and energy of the compositions of regular graphs. *Appl. Math. Lett.* **22** 1136–40.
- [13] Ilic A. 2010. Distance spectra and distance energy of integral circulant graphs. *Linear Algebra Appl.* **433** 1005–14.
- [14] Bose SS, Nath M and Paul S. Distance spectral radius of graphs with r pendent vertices. *Linear Algebra Appl.* **435** 2828–36.
- [15] Nath M and Paul S. 2012. On the distance spectral radius of bipartite graphs. *Linear Algebra Appl.* **436** 1285–96.
- [16] Yu G, Jia H, Zhang H and Shu J. 2012. Some graft transformations and its applications on the distance spectral radius of a graph. *Appl. Math. Lett.* **25** 315–9.
- [17] Lin H, Hong Y, Wang J and Shu J. 2013. On the distance spectrum of graphs. *Linear Algebra Appl.* **439** 1662–9.
- [18] Wang Y and Zhou B. 2013. On distance spectral radius of graphs. *Linear Algebra Appl.* **438** 3490–503.
- [19] Jin Y and Zhang X. 2014. Complete multipartite graphs are determined by their distance spectra ☆. *Linear Algebra Appl.* **448** 285–91.

- [20] Indulal G and Stevanovi D. 2015. The distance spectrum of corona and cluster of two graphs ☆. *AKCE Int. J. Graphs Comb.* **12** 186–92.
- [21] Indulal G and Balakrishnan R. 2016. Distance spectrum of Indu–Bala product of graphs. *AKCE Int. J. Graphs Comb.* **13** 230–4.
- [22] Gutman I. 1978. The energy of a graph. *Ber. Math-Statist. Sect. Forschungszentrum Graz* **103** 1–22.
- [23] Jooyandeh M, Kiani D and Mirzakhah M. 2009. Incidence energy of a graph. *MATCH Commun. Math. Comput. Chem.* **62** 561–72.
- [24] Shi L and Wang H. 2013. The Laplacian incidence energy of graphs. *Linear Algebra Appl.* **439** 4056–62.
- [25] Adiga C and Smitha M. 2009. On maximum degree energy of a graph. *Int. J. Contemp. Math. Sci.* **4** 385–96.
- [26] Gutman I and Wagner S. 2012. The matching energy of a graph. *Discret. Appl. Math.* **160** 2177–87.
- [27] Güngör DA and Çevik AS. 2010. On the Harary energy and Harary Estrada index of a graph. *MATCH Commun. Math. Comput. Chem.* **64** 281–96.
- [28] Adiga C, Sampathkumar E, Sriraj MA and Shrikanth AS. 2013. Color energy of a graph. *Proc. Jangjeon Math. Soc.* **16** 335–51.
- [29] Indulal G, Gutman I and Vijayakumar A. 2008. On distance energy of graphs. *MATCH Commun. Math. Comput. Chem.* **60** 461–72.
- [30] Ramane HS, Gutman I and Revankar DS. 2008. Distance equienergetic graphs. *MATCH Commun. Math. Comput. Chem.* **60** 473–84.
- [31] Ramane HS, Revankar DS, Gutman I, Rao SB, Acharya BD and Walikar HB. 2008. Bound for the distance energy of a graph. *Kragujev. J. Math.* **31** 59–68.
- [32] Zhou B and Ilic A. 2010. On distance spectral radius and distance energy of graphs. *MATCH Commun. Math. Comput. Chem.* **64** 261–80.
- [33] Bozkurt B, Gungor AD and Zhou B. 2010. Note on the distance energy of graphs. *MATCH Commun. Math. Comput. Chem.* **64** 129–34.
- [34] Murugesan N and Meenakshi K. 2011. Energy, distance energy of connected graphs and skew energy pf connected digraphs with respect to its spanning trees and its chords. *Int. J. Comb. Graph Theory Appl.* **4** 77–87.
- [35] Burcu Bozkurt Ş, Adiga C and Bozkurt D. 2013. Bounds on the distance energy and the distance Estrada index of strongly quotient graphs. *J. Appl. Math.* 1–6.
- [36] Zhang X. 2014. The inertia and energy of distance matrices of complete k-partite graphs. *Linear Algebra Appl.* **450** 108–20.
- [37] Zhang X. 2014. The inertia and energy of distance matrices of complete k -partite graphs. *Linear Algebra Appl.* **450** 108–20.
- [38] Lin H, Liu R and Lu X. 2015. The inertia and energy of the distance matrix of a connected graph. *Linear Algebra Appl.* **467** 29–39.
- [39] Heydemann M-C. 1997. Cayley graphs and interconnection networks. In: *Graph Symmetry*. Dordrecht: Springer. 167–224.
- [40] Kandasamy WBV and Smarandache F. 2009. *Groups as graphs*. Judetul Olt, Romania: Editura CuArt.
- [41] Vahidi J and Talebi AA. 2010. The commuting graphs on groups  $D_{2n}$  and  $Q_n$ . *J. Math. Comput. Sci.* **1** 123–7.
- [42] Erfanian A and Tolue B. 2012. Conjugate graphs of finite groups. *Discret. Math. Algorithms Appl.* **04** 1–8.
- [43] Moghaddamfar AR, Shi WJ, Zhou W and Zokayi AR. 2005. On the noncommuting graph associated with a finite group. *Sib. Math. J.* **46** 325–32.
- [44] Anderson DF, Fasteen J and Lagrange JD. 2012. The subgroup graph of a group. *Arab J. Math.* **1** 17–27.

- [45] Alfuraidan MR and Zakariya YF. 2017. Inverse graphs associated with finite groups. *Electron. J. Graph Theory Appl.* **5** 142–54.
- [46] Kakeri F and Erfanian A. 2015. The complement of subgroup graph of a group. *J. Prime Res. Math.* **11** 55–60.
- [47] Abdussakir. 2017. Detour energy of complement of subgroup graph of dihedral group. *Zero-Jurnal Sains, Mat. dan Terap.* **1** 41–8.
- [48] Chartrand G, Lesniak L and Zhang P. 2008. *Graphs and digraphs. 6th ed.* Florida: CRC Press.
- [49] Bondy JA and Murty USR. 2008. *Graph theory.* New York: Springer.
- [50] Abdussakir, Azizah NN and Nofandika FF. 2009. *Teori graf: Topik dasar untuk tugas akhir/skripsi.* Malang: UIN Malang Press.
- [51] Yin S. 2008. Investigation on spectrum of the adjacency matrix and Laplacian matrix of graph  $G_l$ . *WSEAS Trans. Syst.* **7** 362–72.
- [52] Caporossi G, Chasset E and Furtula B. 2009. Some conjectures and properties on distance energy. *Les Cah. du GERAD G–2009–64.*
- [53] Stevanovic D, Milosevic M, Hic P and Pokorny M. 2013. Proof of a conjecture on distance energy of complete multipartite graphs. *MATCH Commun. Math. Comput. Chem.* **70** 157–62.